

**Vyndamax**<sup>™</sup>  
(*tafamidis*)  
61 mg capsules



**MORE TIME FOR YOUR HEART.**  
Live longer. It's possible.

VYNDAMAX is the first and only once-daily, one-capsule treatment for transthyretin amyloid cardiomyopathy (ATTR-CM), a life-threatening disease.

## INDICATION AND IMPORTANT SAFETY INFORMATION

### What are VYNDAQEL and VYNDAMAX?

VYNDAQEL and VYNDAMAX are prescription medicines used to treat adults with cardiomyopathy of wild-type or hereditary transthyretin-mediated amyloidosis (ATTR-CM) to reduce death and hospitalization related to heart problems. It is not known if VYNDAQEL and VYNDAMAX are safe and effective in children.

**Before taking VYNDAQEL or VYNDAMAX, tell your healthcare provider about all your medical conditions, including if you:**

- have liver problems.
- are pregnant or plan to become pregnant.
- are breastfeeding or plan to breastfeed.

Tell your healthcare provider about all the medicines you take including any prescription or over-the-counter medicines, vitamins, and herbal supplements.

**There were no known side effects that happened during treatment with VYNDAQEL or VYNDAMAX.**

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit [www.fda.gov/medwatch](http://www.fda.gov/medwatch), or call 1-800-FDA-1088.

[Click here](#) for the Full Prescribing Information and Patient Information or visit [VYNDAMAX.com](http://VYNDAMAX.com)

# WHAT IS VYNDAMAX?

VYNDAMAX is a transthyretin stabilizer approved to reduce death and hospitalizations related to heart problems in adults with ATTR-CM.

The clinical studies of VYNDAQEL supported the approval of VYNDAMAX.



**Take FOUR 20 mg capsules\*  
ONCE DAILY**



**Take only ONE  
61 mg capsule\*  
ONCE DAILY**

\*Capsules shown are not actual size.



VYNDAQEL is currently available as four 20 mg capsules, taken once a day. VYNDAMAX 61 mg is a once daily, single capsule form of VYNDAQEL that was developed for patient convenience.

VYNDAMAX 61 mg was shown to work the same way as VYNDAQEL 80 mg.

## Selected Safety Information

Before taking VYNDAQEL or VYNDAMAX, tell your healthcare provider about all your medical conditions, including if you have liver problems, or are pregnant or plan to become pregnant, or are breastfeeding or plan to breastfeed.



# VYNDAMAX CAN HELP THOSE WITH ATTR-CM LIVE LONGER WITH FEWER HOSPITAL VISITS.

## • WHAT WOULD YOU DO WITH MORE TIME? •

**In a clinical trial, VYNDAMAX was proven to significantly reduce death and the number of hospital visits related to heart problems.**

After 30 months, the survival rate was higher in the VYNDAMAX group than in the placebo group. People taking VYNDAMAX also saw better results versus those on placebo in measures of quality of life including physical abilities and social activities. These quality-of-life results were noticed in as little as 6 months, and remained consistent throughout the 30-month study.

## HOW VYNDAMAX HELPS FIGHT ATTR-CM

VYNDAMAX binds to the transthyretin protein that causes ATTR-CM, slowing its breakdown and the buildup of dangerous deposits. By doing so, VYNDAMAX can slow the progression of the disease.

It must be taken every day, as prescribed, to increase your chances of staying out of the hospital and living longer.

**Please see Important Safety Information on [page 5](#).**

**[Click here](#) for the Full Prescribing Information and Patient Information or visit [VYNDAMAX.com](http://VYNDAMAX.com)**

# EASY-TO-TAKE VYNDAMAX

## IS THE FIRST AND ONLY ONCE-DAILY, ONE-CAPSULE TREATMENT FOR ATTR-CM.

Patients taking VYNDAMAX experienced no side effects from the medication, and required no additional lab testing or monitoring. VYNDAMAX is a single 61-mg capsule you need to take once daily and works in the same way as four 20-mg capsules of VYNDAQEL.



VYNDAMAX capsules should be swallowed whole and not crushed or cut. If you miss a dose, take it as soon as you remember. If it is almost time for your next dose, skip the missed dose and take the next dose at your regularly scheduled time. Do not take 2 doses at the same time.

### Selected Safety Information

Tell your healthcare provider about all the medicines you take including any prescription or over-the-counter medicines, vitamins, and herbal supplements.


There were no known side effects that happened during treatment with VYNDAMAX and VYNDAMAX.

**Please see Important Safety Information on page 5.**

**[Click here](#) for the Full Prescribing Information and Patient Information or visit [VYNDAMAX.com](http://VYNDAMAX.com)**



**If you have been prescribed VYNDAMAX or VYNDAQEL, VyndaLink can help you:**



Explore financial assistance options including co-pay assistance, alternate funding, or free medicine, if eligible



Coordinate with your doctor and insurance to get VYNDAMAX or VYNDAQEL through a specialty pharmacy



Connect with a Patient Support Navigator for additional support on your treatment journey\*

**For more information visit [VyndaLink.com](https://VyndaLink.com)**

\*If you are interested, you must opt in to this offering. Some offerings are provided through third-party organizations that operate independently and are not controlled by Pfizer. Availability of patient support and eligibility requirements are determined solely by these organizations.

## INDICATION AND IMPORTANT SAFETY INFORMATION

### What are VYNDAQEL and VYNDAMAX?

VYNDAQEL and VYNDAMAX are prescription medicines used to treat adults with cardiomyopathy of wild-type or hereditary transthyretin-mediated amyloidosis (ATTR-CM) to reduce death and hospitalization related to heart problems. It is not known if VYNDAQEL and VYNDAMAX are safe and effective in children.

**Before taking VYNDAQEL or VYNDAMAX, tell your healthcare provider about all your medical conditions, including if you:**

- ☐ have liver problems.
- ☐ are pregnant or plan to become pregnant.
- ☐ are breastfeeding or plan to breastfeed.

Tell your healthcare provider about all the medicines you take including any prescription or over-the-counter medicines, vitamins, and herbal supplements.

**There were no known side effects that happened during treatment with VYNDAQEL and VYNDAMAX.**



## WHAT SHOULD I ASK MY CARDIOLOGIST?



- ☐ If I have ATTR-CM, could VYNDAMAX be right for me?
- ☐ Could VYNDAMAX help slow the progression of my ATTR-CM?
- ☐ What are the possible side effects?
- ☐ How do I take VYNDAMAX?
- ☐ How can I get VYNDAMAX?

Talk to your cardiologist about VYNDAMAX today.

**Vyndamax**<sup>™</sup>  
*(tafamidis)*  
61 mg capsules



Talk to your cardiologist about VYNDAMAX today.

For more information, visit [VYNDAMAX.com](https://www.vyndamax.com)  
or call 1-877-27-VYNDA (1-877-278-9632).

Please see Important Safety Information on [page 5](#).  
[Click here](#) for the Full Prescribing Information and Patient Information  
or visit [VYNDAMAX.com](https://www.vyndamax.com)



## HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use VYNDALCEL and VYNDAMAX safely and effectively. See full prescribing information for VYNDALCEL and VYNDAMAX.

VYNDALCEL® (tafamidis meglumine) capsules, for oral administration  
Initial U.S. Approval: 2019

VYNDAMAX™ (tafamidis) capsules, for oral administration  
Initial U.S. Approval: 2019

### INDICATIONS AND USAGE

VYNDALCEL and VYNDAMAX are transthyretin stabilizers indicated for the treatment of the cardiomyopathy of wild-type or hereditary transthyretin-mediated amyloidosis in adults to reduce cardiovascular mortality and cardiovascular-related hospitalization. (1)

### DOSAGE AND ADMINISTRATION

The recommended dosage is either:

- VYNDALCEL 80 mg orally once daily, or
- VYNDAMAX 61 mg orally once daily (2.1)

- VYNDAMAX and VYNDALCEL are not substitutable on a per mg basis. (2.1)

### DOSAGE FORMS AND STRENGTHS

Capsules: Tafamidis meglumine 20 mg and tafamidis 61 mg. (3)

### CONTRAINDICATIONS

None. (4)

### ADVERSE REACTIONS

To report SUSPECTED ADVERSE REACTIONS, contact Pfizer Inc. at 1-800-438-1985 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).

### USE IN SPECIFIC POPULATIONS

- Pregnancy: Based on animal studies, may cause fetal harm. (8.1)
- Lactation: Advise not to breastfeed. (8.2)

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.

Revised: 04/2020

## FULL PRESCRIBING INFORMATION: CONTENTS\*

- 1 INDICATIONS AND USAGE
- 2 DOSAGE AND ADMINISTRATION
  - 2.1 Recommended Dosage
  - 2.2 Administration Instructions
- 3 DOSAGE FORMS AND STRENGTHS
- 4 CONTRAINDICATIONS
- 6 ADVERSE REACTIONS
  - 6.1 Clinical Trials Experience
- 7 DRUG INTERACTIONS
  - 7.1 BCRP Substrates
- 8 USE IN SPECIFIC POPULATIONS
  - 8.1 Pregnancy
  - 8.2 Lactation
  - 8.3 Females and Males of Reproductive Potential
  - 8.4 Pediatric Use

- 8.5 Geriatric Use
- 10 OVERDOSAGE
- 11 DESCRIPTION
- 12 CLINICAL PHARMACOLOGY
  - 12.1 Mechanism of Action
  - 12.2 Pharmacodynamics
  - 12.3 Pharmacokinetics
- 13 NONCLINICAL TOXICOLOGY
  - 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility
- 14 CLINICAL STUDIES
- 16 HOW SUPPLIED/STORAGE AND HANDLING
- 17 PATIENT COUNSELING INFORMATION

\*Sections or subsections omitted from the full prescribing information are not listed.

## **FULL PRESCRIBING INFORMATION**

### **1. INDICATIONS AND USAGE**

VYNDAQEL and VYNDAMAX are indicated for the treatment of the cardiomyopathy of wild-type or hereditary transthyretin-mediated amyloidosis (ATTR-CM) in adults to reduce cardiovascular mortality and cardiovascular-related hospitalization.

### **2. DOSAGE AND ADMINISTRATION**

#### **2.1 Recommended Dosage**

The recommended dosage is either VYNDAQEL 80 mg (four 20-mg tafamidis meglumine capsules) orally once daily or VYNDAMAX 61 mg (one 61-mg tafamidis capsule) orally once daily.

VYNDAMAX and VYNDAQEL are not substitutable on a per mg basis [*see Clinical Pharmacology (12.3)*].

#### **2.2 Administration Instructions**

The capsules should be swallowed whole and not crushed or cut.

If a dose is missed, instruct patients to take the dose as soon as remembered or to skip the missed dose and take the next dose at the regularly scheduled time. Do not double the dose.

### **3. DOSAGE FORMS AND STRENGTHS**

VYNDAQEL is available as:

- tafamidis meglumine 20 mg: yellow, opaque, oblong capsule, printed with “VYN 20” in red.

VYNDAMAX is available as:

- tafamidis 61 mg: reddish brown, opaque, oblong capsule, printed with “VYN 61” in white.

### **4. CONTRAINDICATIONS**

None.

### **6. ADVERSE REACTIONS**

#### **6.1 Clinical Trials Experience**

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

The data reflect exposure of 377 ATTR-CM patients to 20 mg or 80 mg (administered as four 20-mg capsules) of VYNDAQEL administered daily for an average of 24.5 months (ranging from 1 day to 111 months).

Adverse events were assessed from ATTR-CM clinical trials with VYNDAQEL, primarily a 30-month placebo-controlled trial [*see Clinical Studies (14)*]. The frequency of adverse events in patients treated with VYNDAQEL 20 mg (n=88) or 80 mg (n=176; administered as four 20-mg capsules) was similar to that with placebo (n=177).

In the 30-month placebo-controlled trial, similar proportions of VYNDAQEL-treated patients and placebo-treated patients discontinued the study drug because of an adverse event: 12 (7%), 5 (6%), and 11 (6%) from the VYNDAQEL 80-mg, VYNDAQEL 20-mg, and placebo groups, respectively.

## **7. DRUG INTERACTIONS**

### **7.1 BCRP Substrates**

Tafamidis inhibits breast cancer resistant protein (BCRP) in vitro and may increase exposure of substrates of this transporter (e.g., methotrexate, rosuvastatin, imatinib) following VYNDAQEL 80 mg or VYNDAMAX 61 mg. Dose adjustment may be needed for these substrates.

## **8. USE IN SPECIFIC POPULATIONS**

### **8.1 Pregnancy**

#### Risk Summary

Based on findings from animal studies, VYNDAQEL and VYNDAMAX may cause fetal harm when administered to a pregnant woman. However, limited available human data with VYNDAQEL use in pregnant women (at a dose of 20 mg per day) have not identified any drug-associated risks for major birth defects, miscarriage, or adverse maternal or fetal outcomes. In animal reproductive studies, oral administration of tafamidis meglumine to pregnant rabbits during organogenesis resulted in adverse effects on development (embryofetal mortality, fetal body weight reduction and fetal malformation) at a dosage providing approximately 9 times the human exposure (AUC) at the maximum recommended human dose (MRHD) of VYNDAQEL (80 mg), and increased incidence of fetal skeletal variation at a dosage providing equivalent human exposure (AUC) at the MRHD. Postnatal mortality, growth retardation, and impaired learning and memory were observed in offspring of pregnant rats administered tafamidis meglumine during gestation and lactation at a dosage approximately 2 times the MRHD based on body surface area ( $\text{mg}/\text{m}^2$ ) (*see Data*). Advise pregnant women of the potential risk to a fetus. Report pregnancies to the Pfizer reporting line at 1-800-438-1985.

The estimated background risk of major birth defects and miscarriage for the indicated population is unknown. All pregnancies have a background risk of birth defects, loss, or other adverse outcomes. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2% to 4% and 15% to 20%, respectively.

#### Data

##### *Animal Data*

In pregnant rats, oral administration of tafamidis meglumine (0, 15, 30, and 45 mg/kg/day) throughout organogenesis resulted in decreased fetal body weights at  $\geq 30$  mg/kg/day (approximately 10 times the human exposure at the MRHD based on AUC). The no-observed-adverse-effect-level (NOAEL) for embryofetal development in rats was 15 mg/kg/day (approximately 7 times the human exposure at the MRHD based on AUC).

In pregnant rabbits, oral administration of tafamidis meglumine (0, 0.5, 2, and 8 mg/kg/day) throughout organogenesis resulted in increased embryofetal mortality, reduced fetal body weights, and an increased incidence of fetal malformations at 8 mg/kg/day (approximately 9 times the human exposure at the MRHD based on AUC), which was also maternally toxic. Increased incidences of fetal skeletal variations were observed at doses  $\geq 0.5$  mg/kg/day (approximately equivalent to the human exposure at the MRHD based on AUC).

In the pre- and postnatal study, pregnant rats received oral administration of tafamidis meglumine at doses of 0, 5, 15, or 30 mg/kg/day throughout pregnancy and lactation (Gestation Day 7 to Lactation Day 20). Decreased survival and body weights, delayed male sexual maturation and neurobehavioral effects (learning and memory impairment) were observed in the offspring of dams treated at 15 mg/kg/day (approximately 2 times the MRHD on a mg/m<sup>2</sup> basis). The NOAEL for pre- and postnatal development in rats was 5 mg/kg/day (approximately equivalent to the MRHD on a mg/m<sup>2</sup> basis).

## **8.2 Lactation**

### Risk Summary

There are no available data on the presence of tafamidis in human milk, the effect on the breastfed infant, or the effect on milk production. Tafamidis is present in rat milk (*see Data*). When a drug is present in animal milk, it is likely the drug will be present in human milk. Based on findings from animal studies which suggest the potential for serious adverse reactions in the breastfed infant, advise patients that breastfeeding is not recommended during treatment with VYND AQEL or VYNDAMAX.

### Data

Pregnant and lactating female rats were administered repeated daily oral doses of tafamidis meglumine (15 mg/kg/day) followed by a single oral gavage dose of <sup>14</sup>C-tafamidis meglumine on Lactation Day 4 or 12. Radioactivity was observed in milk by 1 hour post-dose and increased thereafter. The ratio of the highest radioactivity associated with <sup>14</sup>C tafamidis meglumine in milk (8 hours post-dose) vs. plasma (1 hour post-dose) was approximately 1.6 on Day 12, indicating tafamidis meglumine is transferred to milk after oral administration.

## **8.3 Females and Males of Reproductive Potential**

### Contraception

#### *Females*

Based on findings from animal studies, VYND AQEL and VYNDAMAX may cause fetal harm when administered to a pregnant woman [*see Use in Specific Populations (8.1)*]. Consider pregnancy planning and prevention for females of reproductive potential.

## **8.4 Pediatric Use**

The safety and effectiveness of VYND AQEL and VYNDAMAX have not been established in pediatric patients.

## **8.5 Geriatric Use**

No dosage adjustment is required for elderly patients (≥65 years) [*see Clinical Pharmacology (12.3)*]. Of the total number of patients in the clinical study (n=441), 90.5% were 65 and over, with a median age of 75 years.


## **10. OVERDOSAGE**

There is minimal clinical experience with overdose. During clinical trials, two patients accidentally ingested a single VYND AQEL dose of 160 mg without adverse events. The highest dose of tafamidis meglumine given to healthy volunteers in a clinical trial was 480 mg as a single dose. There was one reported adverse event of mild hordeolum at this dose.

## 11. DESCRIPTION


VYNDAQEL (tafamidis meglumine) and VYNDAMAX (tafamidis) contain tafamidis as the active moiety, which is a selective stabilizer of transthyretin.

The chemical name of tafamidis meglumine is 2-(3,5-dichlorophenyl)-1,3-benzoxazole-6-carboxylic acid mono (1-deoxy-1-methylamino-D-glucitol). The molecular formula is  $C_{14}H_7Cl_2NO_3 \cdot C_7H_{17}NO_5$ , and the molecular weight is 503.33 g/mol. The structural formula is:



Tafamidis meglumine 20-mg soft gelatin capsule for oral use contains a white to pink colored suspension of tafamidis meglumine 20 mg (equivalent to 12.2 mg of tafamidis free acid), and the following inactive ingredients: ammonium hydroxide 28%, brilliant blue FCF, carmine, gelatin, glycerin, iron oxide (yellow), polyethylene glycol 400, polysorbate 80, polyvinyl acetate phthalate, propylene glycol, sorbitan monooleate, sorbitol, and titanium dioxide.

The chemical name of tafamidis is 2-(3,5-dichlorophenyl)-1,3-benzoxazole-6-carboxylic acid. The molecular formula is  $C_{14}H_7Cl_2NO_3$ , and the molecular weight is 308.12 g/mol. The structural formula is:



Tafamidis 61-mg soft gelatin capsule for oral use contains a white to pink colored suspension of tafamidis 61 mg and the following inactive ingredients: ammonium hydroxide 28%, butylated hydroxytoluene, gelatin, glycerin, iron oxide (red), polyethylene glycol 400, polysorbate 20, povidone (K-value 90), polyvinyl acetate phthalate, propylene glycol, sorbitol, and titanium dioxide.

## 12. CLINICAL PHARMACOLOGY

### 12.1 Mechanism of Action

Tafamidis is a selective stabilizer of TTR. Tafamidis binds to TTR at the thyroxine binding sites, stabilizing the tetramer and slowing dissociation into monomers, the rate-limiting step in the amyloidogenic process.

## 12.2 Pharmacodynamics

A proprietary TTR stabilization assay was utilized as a pharmacodynamic marker and assessed the stability of the TTR tetramer ex vivo. The TTR stabilization assay quantifies immunoturbidimetric measurement of the stable TTR tetramer in plasma pre- and post-treatment with 2-day in vitro denaturation with urea. Using this proprietary assay, a dose-dependent trend for greater TTR tetramer stabilization is observed for VYNDAREL 80-mg compared to VYNDAREL 20-mg. However, the clinical relevance of a higher TTR tetramer stabilization towards cardiovascular outcomes is not known.

VYNDAREL stabilized both the wild-type TTR tetramer and the tetramers of 14 TTR variants tested clinically after once-daily dosing. Tafamidis also stabilized the TTR tetramer for 25 variants tested ex vivo.

VYNDAREL and VYNDAMAX may decrease serum concentrations of total thyroxine, without an accompanying change in thyroid stimulating hormone (TSH). This reduction in total thyroxine values is probably the result of reduced thyroxine binding to or displacement from transthyretin (TTR) due to the high binding affinity of tafamidis to the TTR thyroxine receptor. No corresponding clinical findings consistent with hypothyroidism have been observed.

Biomarkers associated with heart failure (NT-proBNP and Troponin I) favored VYNDAREL over placebo.

### Cardiac Electrophysiology

At approximately 2.2 times the steady state peak plasma concentration ( $C_{max}$ ) at the recommended dose, tafamidis does not prolong the QTc interval to any clinically relevant extent.

## 12.3 Pharmacokinetics

No clinically significant differences in steady state  $C_{max}$  and area under the plasma concentration over time curve (AUC) of tafamidis were observed for VYNDAMAX 61-mg capsule compared to VYNDAREL administered as four 20-mg capsules.

Tafamidis exposure increases proportionally over single (up to 480 mg) or multiple (up to 80 mg) (1 to 6 times the approved recommended dosage) once daily dosing.

The apparent clearance were similar after single and repeated administration of VYNDAREL 80 mg.

### Absorption

Median tafamidis peak concentrations occurred within 4 hours following dosing.

### *Effect of Food*

No clinically significant differences in the pharmacokinetics of tafamidis were observed following administration of a high fat, high calorie meal.

### Distribution

The apparent steady state volume of distribution of tafamidis meglumine is 16 liters and 18.5 liters for tafamidis. Plasma protein binding of tafamidis is >99% in vitro. Tafamidis primarily binds to TTR.

### Elimination

The mean half-life of tafamidis is approximately 49 hours. The apparent oral clearance of tafamidis meglumine is 0.228 L/h (0.263 L/h for tafamidis). The degree of drug accumulation at steady state after repeated tafamidis daily dosing is approximately 2.5-fold greater than that observed after a single dose.

## *Metabolism*

The metabolism of tafamidis has not been fully characterized. However, glucuronidation has been observed.

## *Excretion*

After a single oral dose of tafamidis meglumine 20 mg, approximately 59% of the dose was recovered in feces (mostly as the unchanged drug) and approximately 22% of the dose was recovered in urine (mostly as the glucuronide metabolite).

## Specific Populations

No clinically significant differences in the pharmacokinetics of tafamidis were observed based on age, race/ethnicity (Caucasian and Japanese) or renal impairment.

## *Patients with Hepatic Impairment*

Patients with moderate hepatic impairment (Child-Pugh Score of 7 to 9) had decreased systemic exposure (approximately 40%) and increased clearance (approximately 68%) of tafamidis compared to healthy subjects. As TTR levels are lower in subjects with moderate hepatic impairment than in healthy subjects, the exposure of tafamidis relative to the amount of TTR is sufficient to maintain stabilization of the TTR tetramer in these patients. No clinically significant differences in the pharmacokinetics of tafamidis were observed in patients with mild hepatic impairment (Child Pugh Score of 5 to 6) compared to healthy subjects. The effect of severe hepatic impairment on tafamidis is unknown.

## Drug Interaction Studies

### *Clinical Studies*

No clinically significant differences in the pharmacokinetics of midazolam (a CYP3A4 substrate) or on the formation of its active metabolite (1-hydroxymidazolam) were observed when a single 7.5-mg dose of midazolam was administered prior to and after a 14-day regimen of VYNDALUR 20-mg once daily.

### *In Vitro Studies*

Cytochrome P450 Enzymes: Tafamidis induces CYP2B6 and CYP3A4 and does not induce CYP1A2. Tafamidis does not inhibit CYP1A2, CYP2B6, CYP2C8, CYP2C9, CYP2C19, CYP3A4/5 or CYP2D6.

UDP glucuronosyltransferase (UGT): Tafamidis inhibits intestinal activities of UGT1A1 but neither induces nor inhibits other UDP glucuronosyltransferase (UGT) systemically.

Transporter Systems: Tafamidis inhibits breast cancer resistant protein (BCRP). In vitro studies and model predictions show that tafamidis has a low potential to inhibit organic anion transporters OAT1 and OAT3 at clinically relevant concentrations. Tafamidis did not show a potential to inhibit Multi-Drug Resistant Protein (MDR1) (also known as P-glycoprotein; P-gp), organic cation transporter OCT2, multidrug and toxin extrusion transporters MATE1 and MATE2K and, organic anion transporting polypeptide OATP1B1 and OATP1B3.

## **13. NONCLINICAL TOXICOLOGY**

### **13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility**

#### Carcinogenesis

There was no evidence of an increased incidence of neoplasia in the transgenic (Tg)-rasH2 mouse following repeated daily administration for 26 weeks at daily doses of 0, 10, 30 or 90 mg/kg. There was no evidence of increased incidence of neoplasia in a 2-year carcinogenicity study in rats at exposures up to 18 times the AUC at the MRHD.

## Mutagenesis

There was no evidence of mutagenicity or clastogenicity in vitro, and an in vivo rat micronucleus study was negative.

## Impairment of Fertility

There were no effects of tafamidis meglumine on fertility, reproductive performance, or mating behavior in the rat at any dose. Rats were dosed daily (0, 5, 15, and 30 mg/kg/day) prior to cohabitation (for at least 15 days for females and 28 days for males), throughout the cohabitation period to the day prior to termination of males and through to implantation of females (Gestation Day 7). No adverse effects were noted on male and female rats in toxicity, fertility, and mating behavior at any dose. The paternal and maternal no observed adverse effect level for reproductive toxicity of tafamidis meglumine is 30 mg/kg/day, approximately 4 times the MRHD on a mg/m<sup>2</sup> basis.

## **14. CLINICAL STUDIES**

Efficacy was demonstrated in a multicenter, international, randomized, double-blind, placebo-controlled study in 441 patients with wild-type or hereditary ATTR-CM (NCT01994889).

Patients were randomized in a 1:2:2 ratio to receive VYNDALCEL 20 mg (n=88), VYNDALCEL 80 mg (administered as four 20-mg VYNDALCEL capsules) (n=176), or matching placebo (n=177) once daily for 30 months, in addition to standard of care (e.g., diuretics). Treatment assignment was stratified by the presence or absence of a variant TTR genotype as well as baseline disease severity (NYHA Class). Transplant patients were excluded from this study. Table 1 describes the patient demographics and baseline characteristics.

**Table 1: Patient Demographics and Baseline Characteristics**

Characteristic	Pooled Tafamidis N=264	Placebo N=177
Age — years		
Mean (standard deviation)	74.5 (7.2)	74.1 (6.7)
Median (minimum, maximum)	75 (46, 88)	74 (51, 89)
Sex — number (%)		
Male	241 (91.3)	157 (88.7)
Female	23 (8.7)	20 (11.3)
TTR Genotype — number (%)		
ATTRm	63 (23.9)	43 (24.3)
ATTRwt	201 (76.1)	134 (75.7)
NYHA Class — number (%)		
NYHA Class I	24 (9.1)	13 (7.3)
NYHA Class II	162 (61.4)	101 (57.1)
NYHA Class III	78 (29.5)	63 (35.6)

Abbreviations: ATTRm = variant transthyretin amyloid, ATTRwt = wild-type transthyretin amyloid

The primary analysis used a hierarchical combination applying the method of Finkelstein-Schoenfeld (F-S) to all-cause mortality and frequency of cardiovascular-related hospitalizations, which was defined as the number of times a subject was hospitalized (i.e., admitted to a hospital) for cardiovascular-related morbidity. The method compared each patient to every other patient within each stratum in a pair-wise manner that proceeded in a hierarchical fashion using all-cause mortality followed by frequency of cardiovascular-related hospitalizations when patients could not be differentiated based on mortality.



This analysis demonstrated a significant reduction ( $p=0.0006$ ) in all-cause mortality and frequency of cardiovascular-related hospitalizations in the pooled VYNDAQEL 20-mg and 80-mg groups versus placebo (Table 2).

**Table 2: Primary Analysis Using Finkelstein-Schoenfeld (F-S) Method of All-Cause Mortality and Frequency of Cardiovascular-Related Hospitalizations**


<b>Primary Analysis</b>	<b>Pooled VYNDAQEL N=264</b>	<b>Placebo N=177</b>
Number (%) of Subjects Alive* at Month 30	186 (70.5)	101 (57.1)
Mean Number of Cardiovascular-related Hospitalizations During 30 months (per patient per year) Among Those Alive at Month 30	0.297	0.455
p-value from F-S Method	0.0006	

\* Heart transplantation and cardiac mechanical assist device implantation are considered indicators of approaching end stage. As such, these subjects are treated in the analysis as equivalent to death. Therefore, such subjects are not included in the count of “Number of Subjects Alive at Month 30” even if such subjects are alive based on 30 month vital status follow-up assessment.

Analysis of the individual components of the primary analysis (all-cause mortality and cardiovascular-related hospitalization) also demonstrated significant reductions for VYNDAQEL versus placebo.

The hazard ratio from the all-cause mortality Cox-proportional hazard model for pooled VYNDAQEL versus placebo was 0.70 (95% confidence interval [CI] 0.51, 0.96), indicating a 30% relative reduction in the risk of death relative to the placebo group ( $p=0.026$ ). Approximately 80% of total deaths were cardiovascular-related in both treatment groups. A Kaplan-Meier plot of time to event all-cause mortality is presented in Figure 1.

**Figure 1: All-Cause Mortality\***



Subjects Remaining at Risk  
(Cumulative events)

Pooled	264	259	252	244	235	222	216	209	200	193	99	0
VYND AQEL	0	5	12	20	29	42	48	55	64	71	78	78
Placebo	177	173	171	163	161	150	141	131	118	113	51	0
	0	4	6	14	16	27	36	46	59	64	75	76

\*Heart transplants and cardiac mechanical assist devices treated as death. Hazard ratio from Cox proportional hazards model with treatment, TTR genotype (variant and wild-type), and NYHA baseline classification (NYHA Classes I and II combined and NYHA Class III) as factors.

There were significantly fewer cardiovascular-related hospitalizations with VYND AQEL compared with placebo with a reduction in risk of 32% corresponding to a Relative Risk Ratio of 0.68 (Table 3).

**Table 3: Cardiovascular-Related Hospitalization Frequency**


	<b>Pooled VYND AQEL N=264</b>	<b>Placebo N=177</b>
Total (%) Number of Subjects with Cardiovascular-related Hospitalizations	138 (52.3)	107 (60.5)
Cardiovascular-related Hospitalizations per Year*	0.48	0.70
Pooled VYND AQEL vs Placebo Treatment Difference (Relative Risk Ratio)*	0.68	
p-value*	<0.0001	

\*This analysis was based on a Poisson regression model with treatment, TTR genotype (variant and wild-type), New York Heart Association (NYHA). Baseline classification (NYHA Classes I and II combined and NYHA Class III), treatment-by-TTR genotype interaction, and treatment-by-NYHA baseline classification interaction terms as factors.

The treatment effects of VYND AQEL on functional capacity and health status were assessed by the 6-Minute Walk Test (6MWT) and the Kansas City Cardiomyopathy Questionnaire-Overall Summary (KCCQ-OS) score,

respectively. A significant treatment effect favoring VYNDAQEL was first observed at Month 6 and remained consistent through Month 30 on both 6MWT distance and KCCQ-OS score (Figure 2 and Table 4).

**Figure 2: Change from Baseline to Month 30 in 6MWT Distance and KCCQ-OS Score**




Abbreviations: 6MWT=6-Minute Walk Test, KCCQ-OS=Kansas City Cardiomyopathy Questionnaire-Overall Summary.

Panel A shows change from Baseline to Month 30 in pooled VYNDAQEL patients compared with placebo patients in 6MWT distance.

Panel B shows change from Baseline to Month 30 in pooled VYNDAQEL patients compared with placebo patients in KCCQ-OS score.

The Kansas City Cardiomyopathy Questionnaire-Overall Summary (KCCQ-OS) score is composed of four domains including Total Symptoms (Symptom Frequency and Symptom Burden), Physical Limitation, Quality of Life, and Social Limitation. The Overall Summary score and domain scores range from 0 to 100, with higher scores representing better health status. All four domains favored pooled VYNDAQEL compared to placebo at Month 30, and demonstrated similar treatment effects to the KCCQ-OS score (Figure 2 and Table 4). The distribution for change from Baseline to Month 30 for KCCQ-OS (Figure 3) shows that the proportion of patients with worse KCCQ-OS scores was lower for the pooled VYNDAQEL-treated group compared to placebo, and the proportion with improved scores was higher (Figure 3).

**Figure 3: Histogram of Change from Baseline to Month 30 in KCCQ-Overall Summary Score**



Abbreviation: KCCQ-OS=Kansas City Cardiomyopathy Questionnaire-Overall Summary.

**Table 4: 6MWT Distance and KCCQ-OS Scores**


Endpoints	Baseline Mean (SD)		Change from Baseline to Month 30, LS Mean (SE)		Treatment Difference from Placebo LS Mean (95% CI)
	Pooled VYND AQEL N=264	Placebo N=177	Pooled VYND AQEL	Placebo	
<b>6MWT (meters)</b>	351 (121)	353 (126)	-55 (5)	-131 (10)	76 (58, 94)
<b>KCCQ-OS</b>	67 (21)	66 (22)	-7 (1)	-21 (2)	14 (9, 18)

Abbreviations: 6MWT = 6-Minute Walk Test; KCCQ-OS = Kansas City Cardiomyopathy Questionnaire-Overall Summary; SD = standard deviation; LS = least squares; SE = standard error; CI = confidence interval

Results from the F-S method represented by win ratio for the combined endpoint and its components (all-cause mortality and frequency of CV-related hospitalization) consistently favored VYND AQEL versus placebo across all subgroups (wild-type, variant and NYHA Class I & II, and III), except for CV-related hospitalization frequency in NYHA Class III (Figure 4). Win ratio is the number of pairs of VYND AQEL-treated patient

“wins” divided by number of pairs of placebo patient “wins.” Analyses of 6MWT and KCCQ-OS also favored VYND AQEL relative to placebo within each subgroup.

**Figure 4: Results by Subgroup, Dose, and Components of Primary Analysis**



Abbreviations: ATTRm = variant transthyretin amyloid, ATTRwt = wild-type transthyretin amyloid, F-S = Finkelstein Schoenfeld, CI = Confidence Interval

\*F-S results presented using win ratio (based on all-cause mortality and frequency of cardiovascular hospitalization)

Heart transplants and cardiac mechanical assist devices treated as death.

Results of the primary analysis, 6MWT at Month 30 and KCCQ-OS at Month 30 were statistically significant for both the 80-mg and 20-mg doses of VYND AQEL vs. placebo, with similar results for both doses.

## 16. HOW SUPPLIED/STORAGE AND HANDLING

VYND AQEL 20-mg (tafamidis meglumine) soft gelatin capsules are yellow, opaque, oblong, and printed with “VYN 20” in red and supplied in the following package configurations:

VYND AQEL Capsules		
Package Configuration	Strength	NDC
Carton of 4 intermediary cartons. Each intermediary carton contains 3 blister cards. Each blister card contains 10 capsules. (120 total capsules)	20 mg	NDC 0069-1975-40

VYNDAMAX 61-mg (tafamidis) soft gelatin capsules are reddish brown, opaque, oblong, and printed with “VYN 61” in white and supplied in the following package configurations:

VYNDAMAX Capsules		
Package Configuration	Strength	NDC
Carton of 3 blister cards. Each blister card contains 10 capsules. (30 capsules total)	61 mg	NDC 0069-8730-30

Store VYND AQEL and VYNDAMAX at controlled room temperature 20°C to 25°C (68°F to 77°F); excursions permitted to 15°C to 30°C (59°F to 86°F) [see USP Controlled Room Temperature].

## 17. PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Patient Information).

### Pregnancy

Report pregnancies to the Pfizer reporting line at 1-800-438-1985. Advise pregnant women and females of reproductive potential of the potential risk to a fetus. Advise females to inform their healthcare provider of a known or suspected pregnancy [*see Use in Specific Populations (8.1)*].

### Lactation

Advise females not to breastfeed during treatment with VYNDAQEL or VYNDAMAX [*see Use in Specific Populations (8.2)*].

### Transthyretin Amyloidosis Outcome Survey (THAOS)

Advise all patients prescribed VYNDAQEL or VYNDAMAX of the availability of the Transthyretin Amyloidosis Outcome Survey (THAOS) registry, that their participation is voluntary, and involves long-term follow-up. THAOS is an international disease registry designed to assess disease progression, genotype/phenotype relationships, and the impact of interventions, including VYNDAQEL and VYNDAMAX on disease progression. For information regarding the registry, visit <https://clinicaltrials.gov>.

This product's label may have been updated. For full prescribing information, please visit [www.pfizer.com](http://www.pfizer.com).



LAB-0497-3.0

## PATIENT INFORMATION

**VYNDALCEL®** (VIN-duh-kel)  
(tafamidis meglumine)  
capsules

**VYNDAMAX™** (VIN-dah-max)  
(tafamidis)  
capsules

### What is VYNDALCEL and VYNDAMAX?

VYNDALCEL and VYNDAMAX are prescription medicines used to treat adults with cardiomyopathy of wild-type or hereditary transthyretin-mediated amyloidosis (ATTR-CM) to reduce death and hospitalization related to heart problems. It is not known if VYNDALCEL and VYNDAMAX are safe and effective in children.

### Before taking VYNDALCEL or VYNDAMAX, tell your healthcare provider about all your medical conditions, including if you:

- have liver problems.
- are pregnant or plan to become pregnant. VYNDALCEL and VYNDAMAX may harm your unborn baby. Tell your healthcare provider right away if you become pregnant or think you may be pregnant during treatment with VYNDALCEL or VYNDAMAX. You may also report your pregnancy by calling the Pfizer reporting line at 1-800-438-1985.
- are breastfeeding or plan to breastfeed. It is not known if VYNDALCEL or VYNDAMAX passes into your breast milk. You should not breastfeed during treatment with VYNDALCEL or VYNDAMAX. Talk to your healthcare provider about the best way to feed your baby during treatment with VYNDALCEL or VYNDAMAX.

Tell your healthcare provider about all the medicines you take including any prescription or over-the-counter medicines, vitamins, and herbal supplements.

### How should I take VYNDALCEL or VYNDAMAX?

- Take **either** VYNDALCEL **or** VYNDAMAX exactly as your healthcare provider tells you to.
- Take **either** VYNDALCEL **or** VYNDAMAX capsule(s) 1 time a day.
- VYNDALCEL or VYNDAMAX capsule(s) should be swallowed whole and not crushed or cut.
- If you miss a dose, take it as soon as you remember. If it is almost time for your next dose, skip the missed dose and take the next dose at your regularly scheduled time. Do not take 2 doses at the same time.
- There is a Transthyretin Amyloidosis Outcome Survey (THAOS) registry for people who receive treatment with VYNDALCEL or VYNDAMAX. Talk to your healthcare provider about how you can take part in this registry. For more information about this registry, go to <https://clinicaltrials.gov>.

### What are the possible side effects of VYNDALCEL and VYNDAMAX?

There were no known side effects that happened during treatment with VYNDALCEL or VYNDAMAX in people with cardiomyopathy of transthyretin-mediated amyloidosis.

You may report side effects to FDA at 1-800-FDA-1088.

### How should I store VYNDALCEL and VYNDAMAX?

- Store VYNDALCEL and VYNDAMAX capsules at room temperature between 68°F to 77°F (20°C to 25°C).
- **Keep VYNDALCEL and VYNDAMAX and all medicines out of the reach of children.**

### General information about the safe and effective use of VYNDALCEL and VYNDAMAX.

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use VYNDALCEL or VYNDAMAX for a condition for which it was not prescribed. Do not give VYNDALCEL or VYNDAMAX to other people, even if they have the same symptoms you have. It may harm them.

You can ask your healthcare provider or pharmacist for information about VYNDALCEL or VYNDAMAX that is written for healthcare professionals.

### What are the ingredients in VYNDALCEL and VYNDAMAX?

#### VYNDALCEL:

**Active ingredient:** tafamidis meglumine

**Inactive ingredients:** ammonium hydroxide 28%, brilliant blue FCF, carmine, gelatin, glycerin, iron oxide (yellow), polyethylene glycol 400, polysorbate 80, polyvinyl acetate phthalate, propylene glycol, sorbitan monooleate, sorbitol, and titanium dioxide

#### VYNDAMAX:

**Active ingredient:** tafamidis

**Inactive ingredients:** ammonium hydroxide 28%, butylated hydroxytoluene, gelatin, glycerin, iron oxide (red), polyethylene glycol 400, polysorbate 20, povidone (K-value 90), polyvinyl acetate phthalate, propylene glycol, sorbitol, and titanium dioxide

LAB-0573-3.0

For more information, go to [www.vyndagel.com](http://www.vyndagel.com) or call 1-800-438-1985.

This Patient Information has been approved by the U.S. Food and Drug Administration

Issued: 04/2020